

On L_1 -distance between first exit times from two regions[‡]

Nikolai Dokuchaev

Department of Mathematics and Computer Sciences,
The University of West Indies, Mona, Kingston 7, Jamaica W.I.

February 1, 2008

Abstract

First exit times from regions and their dependence on variations of boundaries are discussed for diffusion processes. The paper presents an estimate of L_1 -distance between exit times from two regions via expectations of exit times.

Abbreviated title: On L_1 -distance between first exit times

It is known that first exit times from a region for smooth solutions of ordinary equations does not depend continuously on variations of the initial data or on the boundary of the region. But first exit times for non-smooth trajectories of diffusion processes have some path-wise regularity with respect to these variations (some related results can be found in author's papers (1987),(1992)). This paper studies path-wise dependence on fluctuations of the boundary for first exit times of diffusion processes. We present an effective estimate of L_1 -distance between exit times from two regions for a diffusion process via expectations of exit times.

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a standard probability space. Consider a n -dimensional diffusion process $y(t)$ such that

$$dy(t) = f(y(t))dt + \beta(y(t))dw(t), \quad t > 0, \quad (1)$$

$$y(0) = a. \quad (2)$$

Here $w(t)$ is a standard d -dimensional Wiener process, f and β are non-random functions with respective values in $\mathbf{R}^n$ and $\mathbf{R}^{n \times d}$. The random vector a with values in $\mathbf{R}^n$ does not depend on $w(\cdot)$.

All vectors and matrices are real, and $\bar{D}$ denotes the closure of a region D .

*MSC 2000 subject classifications. Primary- 60G17, 60G40 ; secondary - 60J50, 60J60, 60J65.

[‡]*Key words:* diffusion processes, first exit times, variable boundaries

Further, for $x \in \mathbf{R}^n$, we denote by $y^x(t)$ the solution of (1) with the initial condition $y(0) = x$.

For a set $\Gamma \in \mathbf{R}^n$, we denote $\tau(\Gamma) \triangleq \inf\{t : y(t) \in \Gamma\}$. This is the first times of achieving D of the processes $y(t)$. Similarly, we denote $\tau^x(\Gamma) \triangleq \inf\{t : y^x(t) \in \Gamma\}$.

Let D_1, D_2 be two bounded regions in $\mathbf{R}^n$. Let $\Gamma_i = \partial D_i$ be the boundary of D_i .

We assume that the boundaries of D_i are C^1 -smooth. Further, we assume that all the components of the functions f, β are continuously differentiable, and $b(x)b(x)^\top \geq cI_n$, where I_n is the unit matrix in $\mathbf{R}^{n \times n}$, and $c > 0$ is a constant.

Theorem 1 *Let $a \in \bar{D}_1 \cap \bar{D}_2$ with probability 1. Then*

$$\mathbf{E}|\tau(\Gamma_1) - \tau(\Gamma_2)| \leq \max\left(\sup_{x \in D_1 \cap \Gamma_2} \mathbf{E}\tau^x(\Gamma_1), \sup_{x \in D_2 \cap \Gamma_1} \mathbf{E}\tau^x(\Gamma_2)\right). \quad (3)$$

Note the theorem is oriented on the case when $D_1 \setminus D_2 \neq \emptyset$ and $D_2 \setminus D_1 \neq \emptyset$. If $D_1 \subset D_2$ or $D_2 \subset D_1$ then the estimation (3) is obvious.

Example. Let $n = d = 1$, $y^a(t) = a + w(t)$, $D_0 = (0, 1)$, $D_\varepsilon = (\varepsilon, 1 + \varepsilon)$, where $\varepsilon \in [0, 1)$. Let $\Gamma_\varepsilon \triangleq \{\varepsilon, 1 + \varepsilon\}$. Let $\tau_\varepsilon^a = \inf\{t : y^a(t) \in \Gamma_\varepsilon\}$. We have that $\mathbf{E}\tau_\varepsilon^x = (x - \varepsilon)(1 + \varepsilon - x)$ for any $x \in D_\varepsilon$ (it can be found similarly (4) and (9) below). By Theorem 1, it follows that

$$\mathbf{E}|\tau_0^a - \tau_\varepsilon^a| \leq \max\left(\mathbf{E}\tau_\varepsilon^0, \mathbf{E}\tau_0^\varepsilon\right) = \varepsilon(1 - \varepsilon).$$

Remark We assumed that the boundaries and coefficients are smooth, the diffusion is non-degenerate, and the regions are bounded. In fact, these conditions can be lifted provided that the right hand part of (3) is finite. For example, a similar theorem can be obtained for first exit times of the process $(y(t), t)$ from cylindrical regions $D_i \times (0, T)$, $i = 1, 2$, $T > 0$, for the case of time dependent coefficients f and β .

Proof of Theorem 1. Let $v_i = v_i(x) : D_i \rightarrow \mathbf{R}$, $i = 1, 2$, be the solutions in D_i of the Dirichlet problems

$$\mathcal{L}v_i = -1, \quad v_i|_{\Gamma_i} = 0. \quad (4)$$

Here the differential operator

$$\mathcal{L} = \sum_{k=1}^n f_k \frac{\partial}{\partial y_k} + \frac{1}{2} \sum_{k,l=1}^n b_{k,l} \frac{\partial^2}{\partial y_k \partial y_l}, \quad (5)$$

where f_k, y_l , and $b_{k,l}$ are the components of the vectors f, y and the matrix $b = \beta\beta^\top$. As is known, the problem (4) has the unique solution that is twice continuously differentiable up to the boundary.

Let e_1 and e_2 be the indicator functions of the random events $\{\tau(\Gamma_1) > \tau(\Gamma_2)\}$ and $\{\tau(\Gamma_2) > \tau(\Gamma_1)\}$ respectively.

Let $\mathcal{F}_t$ be the filtration generated by $w(t)$ and a .

Let $\hat{\tau} \triangleq \tau(\Gamma_1) \wedge \tau(\Gamma_2)$. The random variables e_i are measurable with respect to the σ -algebras $\mathcal{F}_{\hat{\tau}}, \mathcal{F}_{\tau(\Gamma_i)}$, $i = 1, 2$, associated with the Markov times (with respect to the filtration $\mathcal{F}_t$) $\hat{\tau}$ and $\tau(\Gamma_i)$ (see, e.g., Gihman and Skorohod (1975), Chap. 4, §2). Using Itô's formula, we obtain the equality

$$\begin{aligned} \mathbf{E} \{e_1 v_1[y(\tau(\Gamma_2))]\} &= -\mathbf{E} \{e_1 \{v_1[y(\tau(\Gamma_1))] - v_1[y(\tau(\Gamma_2))]\}\} \\ &= -\mathbf{E} \left\{ e_1 \int_{\hat{\tau}}^{\tau(\Gamma_1)} \mathcal{L}_1 v_1[y(t)] dt \right\} \\ &= \mathbf{E} \{e_1 [\tau(\Gamma_1) - \hat{\tau}]\} \\ &= \mathbf{E} \{e_1 [\tau(\Gamma_1) - \tau(\Gamma_2)]\}. \end{aligned} \tag{6}$$

If we replaced the indices 1, 2 in (6) by 2, 1, we get similarly that

$$\mathbf{E} \{e_2 v_2[y(\tau(\Gamma_1))]\} = \mathbf{E} \{e_2 [\tau(\Gamma_2) - \tau(\Gamma_1)]\}. \tag{7}$$

Clearly,

$$\mathbf{E} |\tau(\Gamma_1) - \tau(\Gamma_2)| = \mathbf{E} \{e_1 [\tau(\Gamma_1) - \tau(\Gamma_2)]\} + \mathbf{E} \{e_2 [\tau(\Gamma_2) - \tau(\Gamma_1)]\}. \tag{8}$$

We have that

$$v_i(x) = \mathbf{E} \tau^x(\Gamma_i). \tag{9}$$

Then it follows from (6)-(8) that

$$\begin{aligned} \mathbf{E} |\tau(\Gamma_1) - \tau(\Gamma_2)| &\leq \mathbf{E} \{e_1 [\tau(\Gamma_1) - \tau(\Gamma_2)]\} + \mathbf{E} \{e_2 [\tau(\Gamma_2) - \tau(\Gamma_1)]\} \\ &= \mathbf{E} \{e_1 \{v_1[y(\tau(\Gamma_2))]\}\} + \mathbf{E} \{e_2 \{v_2[y(\tau(\Gamma_1))]\}\} \\ &\leq \max \left(\sup_{x \in D_1 \cap \Gamma_2} v_1(x), \sup_{x \in D_2 \cap \Gamma_1} v_2(x) \right). \end{aligned}$$

Now the assertion of the Theorem 1 follows. $\square$

References

- Dokuchaev, N.G. (1992). The integral estimations for ordinary differential equations with a discontinuity on a domain boundary. *Differential equations* **28**, N 11, 1540-1549.
- Dokuchaev, N.G. (1987) On moment's of first exit for homogeneous processes of diffusion type. *Theory of Probability and its Applications* **31**, N 3, 3, 497-498.
- Gihman, I.I., and Skorohod, A.V. (1975). *The Theory of Stochastic Processes*. Vol. 2. Springer-Verlag, New York.

Nikolai Dokuchaev,
Department of Mathematics and Computer Science, The University of West Indies,
Mona, Kingston 7, Jamaica W.I.
email ndokuch@uwimona.edu.jm